

**NEUTRINO PROPAGATION THROUGH A
FLUCTUATING SUN**C.P. BURGESS* *and* D. MICHAUD*Physics Department, McGill University
3600 University St., Montréal, Québec, Canada, H3A 2T8.***ABSTRACT**

We summarize a general formulation of particle propagation in fluctuating media, as applied to the description of neutrino propagation through the sun. It contains the familiar MSW effective hamiltonian, plus corrections which describe neutrino interactions with fluctuations in the medium. An estimate of the size of these corrections for a simple model of solar fluctuations is made, with the conclusion that they can produce surprisingly large effects since they grow with the correlation length of the fluctuation. For MSW oscillations, the leading effect for small fluctuations is to diminish the quality of the resonance, making the suppression of the ^7Be neutrinos an experimental probe of fluctuations deep within the sun. Fluctuations can also provide an energy-independent suppression factor of $\frac{1}{2}$, away from the resonant region, even for small vacuum mixing angles. To be even potentially detectable, density fluctuations must be correlated on scales of hundreds of kilometres, and have amplitudes of a few percent or larger.

The next generation of solar-neutrino detectors, including SNO and Super Kamiokande,¹ bring us into a new era, in which neutrino solutions to the solar-neutrino problem will be put under detailed experimental scrutiny. It therefore behooves theorists to understand in detail the corrections to the predictions of standard solar models, with and without neutrino oscillations. Since standard treatments² of MSW oscillations replace the solar medium with an average electron density, one such correction consists of how fluctuations in solar properties about this average can affect neutrino propagation. We very briefly review here a method³ for treating such fluctuations systematically. Limitations of space preclude our discussing related approaches^{4,5} to these issues.

*Talk presented to Neutrino 96, Helsinki, Finland, June 1996.

A great deal is known about how particles like photons and neutrons propagate through materials, and general techniques have been developed to describe this propagation. Our approach is based on an adaptation of these ideas to neutrino physics. In order to keep approximations explicit, we start with the general case, and then specialize to neutrinos within the sun. In its most general context we consider two sectors, A and B , where A describes the degrees of freedom we wish to follow – such as neutrino flavour – and B describes the other degrees of freedom, which we do not intend to measure – like those of the solar medium. (There is a generalization of the approach for which a limited measurement to B is also performed.³) We imagine B to be essentially unchanged by its interaction with sector A . All predictions for measurements purely within sector A are described by the reduced density matrix, defined by tracing the full density matrix over the degrees of freedom in B : $\rho_A(t) \equiv \text{Tr}_B[\rho(t)]$.

The time evolution of ρ_A is computed using the hamiltonian $H = H_0 + V$, where $H_0 = H_A + H_B$ describes the separate evolution of sectors A and B , and V describes their mutual interaction. There are two important timescales which govern this time evolution, and considerable simplification arises when they are very different. The first scale, τ_p , is defined as the time beyond which perturbation theory in V fails. The second scale, τ_c , is the correlation time, defined as the time above which $\langle \delta V(t) \delta V(t') \rangle$ becomes negligible (if such a time exists). Here the average, $\langle \cdots \rangle = \text{Tr}_B(\rho \cdots)$, is over sector B , and $\delta V \equiv V - \bar{V}$ is the deviation of the interaction hamiltonian from the effective interaction, $\bar{V}$, which describes the average evolution in sector A . In perturbation theory $\bar{V}(t) = \langle V(t) \rangle - i \int_{t'}^t d\tau \langle \delta V(t) \delta V(\tau) \rangle + O(V^3)$.

If $\tau_c \ll \tau_p$, then ρ_A satisfies a master equation which can be derived in perturbation theory, and then integrated to obtain its large-time behaviour, even for $t > \tau_p$. It is given, to second order in V , by:

$$\begin{aligned} \frac{\partial \rho_A}{\partial t} = & -i [\bar{V}(t) \rho_A(t) - \rho_A(t) \bar{V}^*(t)] \\ & + \int_{t'}^t d\tau \text{Tr}_B (\delta V(t) \rho(t) \delta V(\tau) + \delta V(\tau) \rho(t) \delta V(t)) + O(V^3). \end{aligned} \quad (1)$$

These ideas may be applied to obtain an equation describing the evolution of neutrino flavour within the solar medium.³ The result is a two-by-two matrix equation:

$$\frac{\partial \rho_f}{\partial t} = -i [V_0 + \bar{V}_1, \rho_f] - 2 G_F^2 \mathcal{A} [(g^e)^2 \rho_f + \rho_f (g^e)^2 - 2 g^e \rho_f g^e], \quad (2)$$

where ρ_f is the 2×2 neutrino-sector density matrix in flavour space, $V_0 \approx k + \frac{m^\dagger m}{2k} + \cdots$, where k is the neutrino momentum and m is

its mass matrix. $\bar{V}_1(t) \equiv \sqrt{2} G_F \langle g^e n_e(t) + g^n n_n(t) \rangle$ is the first-order effective interaction, with $g^e = \text{diag}(1, 0)$, and $g^n = \text{diag}(-\frac{1}{2}, -\frac{1}{2})$ being the 2×2 matrices which describe the charged-current (neutral-current) neutrino coupling to the electron (neutron) density, n_e (n_n). Because g^n is proportional to the unit matrix, the neutron density drops out of eq. (2). The first term of eq. (2) gives the usual MSW evolution. All second-order terms are proportional to the coefficient:

$$\mathcal{A}(t) = \int_{t'}^t d\tau \langle \delta n_e(t) \delta n_e(\tau) \rangle, \quad (3)$$

which controls the size of fluctuation effects.

It is instructive to compare the order of magnitude of the various terms in eq. (2). The flavour-changing part of V_0 is $m^\dagger m/k$; while the term involving V_1 is $\sim G_F g^e \langle n_e \rangle$. These two terms are comparable over part of the solar interior, and their interplay gives rise to the MSW resonant oscillation. The contribution of the remaining term is of order $G_F^2 \langle \delta n_e \delta n_e \rangle \tau_c$, which is competitive with the others if $G_F \langle \delta n_e \delta n_e \rangle \tau_c \sim \langle n_e \rangle$. Writing $\langle \delta n_e \delta n_e \rangle = \epsilon^2 \langle n_e \rangle^2$, this condition becomes $\epsilon^2 G_F \langle n_e \rangle \tau_c \sim 1$. Since, deep within the solar interior, $G_F \langle n_e \rangle \lesssim (100 \text{ km})^{-1}$, significant fluctuation effects require $\epsilon^2 \tau_c \gtrsim 100 \text{ km}$.

These estimates are borne out by more detailed calculations.³ When applied to thermal fluctuations, they reproduce the standard result for neutrino scattering from thermally-distributed electrons, giving a negligible scattering rate within the sun. We have identified only one potential source of fluctuations within the sun whose effects may be nonnegligible. These fluctuations arise due to the variation, over space and time, of the mean electron density within the sun. The propagation of any particular neutrino through such variations is perfectly well described by the MSW treatment. In general, however, successive neutrinos do *not* encounter the *same* electron density profile. This could be because different neutrinos are produced at different places within the sun, and so pass through different regions while *en route* to the earth. Alternatively, successive neutrinos produced at the same point within the sun could pass through different electron densities because the density profile itself generally varies in time – such as when helioseismic waves are present. As a result, the neutrino flux to which a detector is exposed can be thought to have experienced an *ensemble* of density profiles, which must be averaged to obtain the integrated neutrino signal as seen by a detector on earth.

We have computed³ how the MSW resonance is altered due to two kinds of ensembles of this type which are of particular interest. These ensembles differ in the nature of the correlations which they assume. One choice assumes fluctuations which are uncorrelated in position

space, *i.e.* $\delta n_e(x)$ is uncorrelated with $\delta n_e(y)$ for sufficiently large $|\mathbf{x}-\mathbf{y}|$. This kind of local fluctuation is typical of many condensed-matter applications, and could arise in the sun from a number of sources. The second choice consists of fluctuations which are uncorrelated in *momentum* space, with $\delta n_e(k)$ uncorrelated with $\delta n_e(k')$, where $k \neq k'$ label an appropriate set of density-oscillation modes. This second kind of ensemble is intended as a simple model of how neutrinos interact with helioseismic waves, since these have been observed to exist in the sun.

It is straightforward to generalize the MSW analysis to arrive at a variation of Parke's formula⁶:



$$P_e(t) = \frac{1}{2} + \left(\frac{1}{2} - P_J\right) \lambda \cos 2\theta_m(t') \cos 2\theta_m(t), \quad (4)$$

in which $P_J = \exp \left[-\frac{\pi}{2} \left(\frac{\sin^2 2\theta_V}{\cos 2\theta_V} \right) \left(\frac{\delta m^2 h}{2k} \right) \right]$ is the usual 'jump' probability as one passes through the resonance point, with h is the scale height for the electron density and θ_V , θ_m and $\delta m^2 = m_h^2 - m_l^2$ are the vacuum and matter mixing angles, and the difference between the squares of the masses, for the two neutrino states. Fluctuation effects enter eq. (4) through the damping factor $\lambda = \exp \left[-2 \int_{t'}^t dx G_F^2 \mathcal{A}(x) \sin^2 2\theta_m(x) \right]$.

Notice that the fluctuations dominantly act to *damp* the neutrino oscillations. This damping is a reflection of the conversion of the incoming neutrino from a pure to a mixed state as it interacts with the fluctuations. The damping ruins the MSW resonance condition, thereby reducing the effectiveness of the flavour change as the neutrino passes through the resonant point. Our numerical results (presented in the figures) show that small fluctuations first become noticeable where the resonant flavour change is the strongest. Since ${}^7\text{Be}$ neutrinos are suppressed by MSW oscillations in this way, a deviation from the MSW prediction for the strength of the ${}^7\text{Be}$ neutrino flux could be evidence for electron density fluctuations deep within the solar interior.

For sufficiently large times, eq. (4) has the universal prediction: $P_e \rightarrow \frac{1}{2}$. This limit is also seen in our numerical integrations, which agree well with eq. (4) throughout parameter space. This suggests a new solution for the solar neutrino problem: an approximately energy-independent suppression of the solar neutrino flux by a factor of 2 due to decoherence by solar fluctuations. Such a suppression would look much like large-angle vacuum oscillations, even though it could arise through fluctuations with small vacuum mixing angles.

The figures present a summary of the results of a numerical integration of eq. (2), using eq. (3) for density fluctuations which are uncorrelated in either position and momentum space.³ As is seen from these figures, deviations from MSW predictions require fluctuations whose amplitude is, at the very least, a few percent.



The ν_e survival probability as a function of the neutrino energy, in units of Δm^2 . Figure 1 (Figure 2) uses fluctuations which are uncorrelated in position (momentum) space. The vacuum mixing angle is taken to be 0.1. *epsilon* denotes the fractional amplitude of the fluctuation while in Figure 1 ℓ represents the correlation length.

Acknowledgements

We would like to thank the Conference organizers for providing such a splendid setting, and for their kind invitation to present our work. Our funds have been provided by the Natural Sciences and Engineering Research Council of Canada and les Fonds pour la Formation de Chercheurs et l'Aide à la Recherche du Québec.

References

1. For a current update, see the contributions of Y. Suzuki and A.B. McDonald, this volume.
2. Some excellent reviews are: J.N. Bahcall, *Neutrino Astrophysics*, Cambridge University Press, 1989; S. Turck-Chièze *et.al.*, *Phys. Rev.* **C230** (1993) 57; W. Haxton, *Ann. Rev. of Astr. and Astrophys.* **33** (1995) 459.
3. C.P. Burgess and D. Michaud, preprint McGill-96/18, hep-ph/9606295.
4. F.N. Loreti and A.B. Balantekin, *Phys. Rev.* **D50** (1994) 4762; F.N. Loreti, Y.Z. Qian, G.M. Fuller and A.B. Balantekin, *Phys. Rev.* **D52** (1995) 6664; A.B. Balantekin, J.M. Fetter and F.N.Loreti, preprint astro-ph/9604061.
5. H. Nunokawa, A. Rossi, V.B. Semikoz and J.W.F. Valle, *Nucl. Phys.* **B472** (1996) 195; A. Rossi, this volume.
6. S.J. Parke, *Phys. Rev. Lett.* **57** (1986) 1275.